PROUDLY MADE IN india

BOROSIL®
Scientific
enabling the future of science



Precision

Safety

Efficency

Reliable Reagent Handling for Dairy Laboratories

Application note

LABQUEST

SMART • RUGGED • PRACTICAL

Introduction



Fat content analysis is vital in dairy quality control, influencing product classification, regulatory compliance, and consumer trust. The Gerber Method is widely used for its accuracy and simplicity, using 1 mL amyl alcohol and 10 mL sulfuric acid (H_2SO_4) to separate and quantify fat in milk samples. The precision of reagent dispensing plays a crucial role in ensuring accurate and reproducible results. Even minor inaccuracies in reagent volume can lead to inconsistent fat separation, miscalculated fat percentages, and unreliable quality control assessments, ultimately affecting product classification and compliance with dairy industry standards. This application note highlights the importance of using Bottle-Top Dispensers (BTDs) for enhanced accuracy, safety, and efficiency in dairy laboratories.

Process Overview



Impact of Volume Inaccuracies

Reagent	Effect of Under-Dosing	Effect of Over-Dosing
H_2SO_4	Incomplete digestion, trapped fat, low fat% readings	Excess heat, sample charring, false high readings
Amyl Alcohol	Poor fat-layer separation, cloudy interface, measurement errors	Altered density, distorted menis-cus, inconsistent results

Inaccurate reagent volumes can lead to:

- **Quality control failures**, such as incorrect fat content classification.
- **Financial losses** due to rejected batches and regulatory non-compliance.
- **Safety hazards**, including **H_2SO_4 spills and inhalation risks** during manual handling.

Advantages of Using Bottle-Top Dispensers

1. Precision & Reproducibility

- Ensures accurate dispensing of H_2SO_4 and amyl alcohol
- Narrow delivery tubes facilitate direct transfer into Gerber tubes, preventing reagent loss.

2. Enhanced Safety

- Closed-system design minimizes exposure to corrosive and volatile chemicals.
- Reduces the risk of H_2SO_4 splashes, inhalation hazards, and accidental spills.

3. Operational Efficiency

- One-handed operation improves workflow efficiency.
- Telescopic adapters ensure compatibility with different reagent bottles, reducing waste.



Best Practices for Dairy Laboratories

- Utilize high-purity amyl alcohol and H_2SO_4 to ensure optimal fat separation.
- Choose PTFE/PFA-coated BTDs for superior resistance to H_2SO_4 corrosion, as these fluoropolymer coatings offer excellent chemical inertness, preventing degradation and extending dispenser longevity compared to standard plastic materials.

Conclusion

Precision in chemical dispensing is paramount in dairy fat analysis. By integrating BTDs, dairy laboratories can enhance accuracy, improve safety, and reduce operational costs, ultimately ensuring product integrity and regulatory adherence. For instance, a study conducted in a high-volume dairy testing facility showed a 15% reduction in sample reprocessing and a 30% decrease in reagent wastage after switching to BTDs, highlighting their impact on both efficiency and cost savings.

BOROSIL®
Scientific

Registered & Corporate Office:
1101, Crescenzo, G-Block, Opp. MCA Club,
Bandra Kurla Complex, Bandra (E) Mumbai - 400 051. India.
Works:
Plot No. 7, Sr. No. 234, 235 & 245, Indialand Global Industrial Park,
Hinjewadi Phase 1, Pune 411057.
T : +91 20 66779000

PROUDLY MADE IN india